

INTRODUCTION

Blood cultures enable the detection of bacteria and/or fungus in the blood and guide the appropriate selection of antimicrobials^{1,2}. Accuracy of test results rely on correct blood volume to improve confirmation of bacteraemia or fungaemia and minimise the risk of contamination.

This guidance is written in line with the NSW Health Policy Directive *Intravascular Access Devices (IVAD) - Infection Prevention and Control (PD2019_040)*. It is intended for **paediatric patients only** (1 month - 16 years). For adult and neonatal blood culture sampling guidance refer to the Clinical Excellence Commission (CEC) [sepsis website](#).

WHEN TO TAKE BLOOD CULTURES IN CHILDREN

Blood cultures are recommended for children with any of the following:

- suspected sepsis as per the CEC [Paediatric Sepsis Pathway](#)
- new confusion/change in behaviour or irritability
- fever (or history of fever) and immunocompromised, including neutropenia
- fever or evidence of infection with a vascular access device or recent surgery
- fever and recent overseas travel

It is worth noting that the time of onset of fever may not correlate with bacteraemia in children. Instead, blood culture collection should be performed when there is a clinical suspicion of sepsis regardless of whether the child has a fever³.

IMPORTANT POINTS TO REMEMBER

- Blood cultures are the 'gold standard' for the detection of microbial pathogens related to bacteraemia and sepsis⁴
- Adequate volume of blood is needed to be able to culture bacteria and fungi^{1,5}
- Always use aseptic technique - correct technique may help reduce the risk of cross contamination and a false positive test result^{4,6}
- Ensure hand hygiene is performed as per the [5 Moments for Hand Hygiene](#)⁶
- In patients with a central venous access device (CVAD) and suspected sepsis, one set of blood cultures should be taken from each lumen of the CVAD as well as one from a peripheral site⁷

BLOOD CULTURE PROCEDURE

1. Inform child/parent/carer of the procedure, explain the purpose, and obtain verbal consent.
2. Prepare child/family/carer using appropriate distraction therapies for the child's age.
3. Perform hand hygiene.
4. Check pathology request form, patient identification and chlorhexidine allergy history.
5. Perform hand hygiene, assemble and prepare the following equipment on a procedure trolley:
 - Alcohol-based hand rub
 - Appropriate blood culture bottles for the child's age/size (aerobic +/- anaerobic bottle) based on the likelihood of anaerobic infection and blood sample volume; check expiry date of each bottle
 - Label each bottle with name, Medical Record Number (MRN), date/time of collection of blood and location of site used for each set; do not cover bar codes or the bottom of the bottle.
 - Gloves or sterile gloves, small dressing pack, tape, tourniquet(s), eye protection
 - Alcohol (ethanol or isopropyl alcohol), or alcohol with chlorhexidine, according to local procedure
 - Winged infusion set with leash and Vacutainer® sleeve designed to fit over the neck of the blood culture bottle; if unavailable use a winged infusion set with luer adapter and syringe
 - Sharps container
6. Put on protective eyewear and perform hand hygiene.
7. Remove the cap of each blood culture bottle and scrub the vial stoppers using alcohol, or alcohol with chlorhexidine, and allow to dry completely.
8. Position child appropriately, apply tourniquet to palpate and identify appropriate vein.
9. Using alcohol, or alcohol with chlorhexidine⁶ disinfect the venepuncture site using a circular 'scrubbing' motion, spiralling out from the planned venepuncture site. Use a fresh swab for each 'scrub'. Perform 2-3 'scrubs' and do this for a total of 1-2 minutes, then allowing the site to dry. After cleaning if re-palpation of the site is expected, use of sterile gloves and sterile procedure is recommended.
10. Perform hand hygiene.
11. Put on gloves and use aseptic non-touch technique. If re-palpation of the venepuncture site occurs, it must be re-cleansed (return to step 8)^{1,4,6}.
12. Perform venepuncture using winged infusion set with luer adapter and Vacutainer® sleeve.
13. Fill each bottle (3 mL in paediatric bottles and 10 mL in adult bottles; the accuracy of the result correlates with the volume of blood collected)². Invert bottles gently several times to prevent clotting.
 - Always collect the blood culture bottles FIRST then, if required, collect additional blood pathology tubes.
 - Release tourniquet, remove needle (with safety sheath applied), tape cotton ball across the skin site and apply pressure (where possible request child/parent/guardian to take over application of pressure).
14. Repeat steps 8 – 13 if collecting a second set of blood cultures from a different peripheral site.
15. Discard sharps, collect all rubbish/dirty items and dispose of appropriately.
16. Remove gloves and perform hand hygiene; remove eye protection and perform hand hygiene.
17. Place bottles into the biohazard bag and arrange to send to the laboratory with the request form. Transport bottles at room temperature.
18. Document in the health record the number of sets of blood cultures that have been taken, the sites and reason for site choice if this differs from a peripheral site.
19. Perform hand hygiene.
20. Explain to child/family/carer/guardian that results may not be available for 48 hours.

FREQUENTLY ASKED QUESTIONS

Why bother taking blood cultures when most results come back negative?

Studies show that insufficient blood sample volume increases the risk of a false negative result^{1,4,5}. The optimal recovery of bacteria and fungi from blood depends on culturing an adequate volume of blood. It is therefore important to follow the procedure for taking blood cultures to collect a sufficient blood sample volume. The direct correlation between the volume of blood cultured and yield relates to the low number of colony forming units in a millilitre of blood. For each additional millilitre of blood cultured, the yield of microorganisms recovered from blood increases⁵.

A positive result provides direct evidence of infection, enabling the antibiotic treatment to be directed against the demonstrated pathogen(s). Furthermore, cumulative antibiograms can be constructed by summarising antibiotic susceptibility of blood isolates which then supports development of reliable empiric antibiotic treatment guidelines. Click [here](#) for National Healthcare Safety Network (NHSN) Centers for Disease Control and Prevention Organism list.

What should I do if only a very small amount of blood is collected?

It is recommended that 0.5 mL to 1 mL of blood is collected for infants, and 1 to >30 mL for older children⁷. This should be placed in an aerobic bottle. The rationale for this is that most bacteraemia is caused by aerobic and facultative bacteria, which will

be recovered better from aerobic bottles. In addition, pathogenic yeasts are recovered almost exclusively from aerobic bottles, as are strict aerobes, such as *Pseudomonas* and *Stenotrophomonas*.

The volume of blood drawn for culture is the most important determinant of the sensitivity of detection of bacteraemia or fungaemia. Where possible, seek expert help in obtaining a larger blood sample.

Can I collect blood cultures from an intravenous cannula?

Collection of blood cultures via an intravenous cannula is NOT the preferred method¹. However, if this method is used it should only be from a cannula that has been freshly inserted and a second specimen should be obtained from a peripheral site^{1,8}.

How are blood cultures stored prior to transport to the laboratory?

Storage should be at room temperature and never refrigerated⁵. Where transport is delayed, the facility should liaise with the receiving laboratory to establish guidance on sample storage.

REFERENCES

1. Cheng MP, Stenstrom R, Paquette K, et al. Blood Culture Results Before and After Antimicrobial Administration in Patients With Severe Manifestations of Sepsis: A Diagnostic Study. *Ann Intern Med* 2019; 171:547.
2. Clinical Excellence Commission. Healthcare Associated Infection (HAI) Clinical Indicator Manual. Version 3.2. December 2020. Available at: https://www.cec.health.nsw.gov.au/_data/assets/pdf_file/0009/258372/HAI-Clinical-indicator-manual.pdf
3. Kee PP, Chinnappan M, Nair A, Yeak D, Chen A, Starr M, Daley AJ, Cheng AC, Burgner D. Diagnostic Yield of Timing Blood Culture Collection Relative to Fever. *Pediatr Infect Dis J*. 2016 Aug;35(8):846-50.
4. Fabre V, Sharara SL, Salinas AB, et al. Does This Patient Need Blood Cultures? A Scoping Review of Indications for Blood Cultures in Adult Nonneutropenic Inpatients. *Clin Infect Dis* 2020; 71:1339.
5. CLSI. Principles and procedures for blood cultures; Approved guideline. CLSI document M47-A. Wayne, PA: Clinical and Laboratory Standards Institute; 2007
6. Infection Prevention and Control Policy (PD2017_013) https://www1.health.nsw.gov.au/pds/ActivePDSDocuments/PD2017_013.pdf
7. Intravascular Access Devices (IVAD) - Infection Prevention & Control (PD2019_040) https://www1.health.nsw.gov.au/pds/ActivePDSDocuments/PD2019_040.pdf
8. Al-Hamad A, Al-Ibrahim M, Alhajhouj E, Al-Alshaikh Jaffer W, Altowaileb J, Alfaraj H. Nurses' competency in drawing blood cultures and educational intervention to reduce the contamination rate. *J Infect Public Health*. 2016;9(1):66-74.