





Name of medicine	Brand names	Used for	Directions	Daily time table				Change	Comments
				 Morning	 Noon	 Evening	 Bedtime		
Perindopril arginine 10mg tablet	Coversyl Prexum	Treating high blood pressure	Take 1 tablet in the morning.	1				unchanged	
Pantoprazole 20mg tablet	Somac Salpraz	Treating heartburn	Take 2 tablets in the morning.	2				increased dose	To be reviewed by GP after 1 week.
Aspirin 100mg tablet	Astrix Cardiprin Cartia	Preventing stroke and heart attack	Take 1 tablet in the morning.	1				unchanged	Take with breakfast
Metformin 500mg tablet	Diabex Diaformin	To control amount of sugar in the blood	Take 1 tablet in the morning 1 tablet at noon And 1 tablet in the evening.	1	1	1		unchanged	Take during or immediately after food
Glyceryl trinitrate 5mg/24 hours transdermal patch	Transiderm- Nitro 25 Nitro-Dur	Treating chest pain	Apply 1 patch in the morning And remove at night.	Apply			Remove	unchanged	Apply to hair free area of chest or upper arm
Paracetamol 500mg tablet	Panadol Panamax	To reduce pain	Take 2 tablets every 4 to 6 hours when needed for pain. Do NOT take more than 8 tablets in a day.	Take: 2 tablets every 4 to 6 hours when needed for pain. Do NOT take more than 8 tablets in a day.				unchanged	

The following medicines were CEASED. Do not take these medicines without further advice.

Name of medicine	Brand names	Date ceased	Explanation
Meloxicam 15mg tablet	Mobic Movalis	10/07/2021	Making heartburn worse

Allergies and adverse drug reactions

Date	Medicine/causal agent	Reaction
	No known drug allergies or adverse drug reactions	

Patient Medication List

Allergies: No known drug allergies or adverse drug reactions	Patient Information: EXAMPLE, Patient MRN: 1234567 Date of Birth: 12/03/1956 Admission Date and Time: 10/07/2021 13:22
--	---

	What	How much	When	Why	Instructions
New	candesartan (Atacand 8 mg oral tablet)	1 tablet(s) taken by mouth	Daily in the morning	IHD (Improve heart function)	Replaces Coversyl (perindopril) tablets
New	paracetamol (Panamax 500 mg oral tablet)	1 tablet(s) taken by mouth	FOUR times a Day as needed for fever/pain		Do not take more than 8 in 24 hours
New	trimETHOPRIM (Alprim 300 mg oral tablet)	1 tablet(s) taken by mouth	Daily at night	Urinary Tract Infection	
Changed - Dose decreased	metoprolol (metoprol 50 mg oral tablet)	1 tablet(s) taken by mouth	TWICE a day	Hypertension (High blood pressure)	Same as Minax
Unchanged	aspirin (Cartia 100 mg oral enteric tablet)	1 tablet(s) taken by mouth	Daily in the morning (with or after food)	Stroke prevention	Swallow whole. Do not crush or chew tablet.
Unchanged	atorvastatin (Lipitor 40 mg oral tablet)	1 tablet(s) taken by mouth	Daily in the morning	High Cholesterol	
Unchanged	tamsulosin (Flomaxtra 0.4 mg oral modified release tablet)	1 tablet(s) taken by mouth	Daily in the morning	Prostate	Swallow whole. Do not crush or chew tablet.

Stopped Medications

	What	How much	When	Comment
Stop taking	perindopril (coVERSYL 5 mg oral tablet)	1 tablet(s) taken by mouth	Daily in the morning	Stopped because of a dry cough

Additional Information: If you have any questions please contact the Pharmacy Department to speak to a pharmacist at MyHospital.on.(0291234567)	Prepared By: One Medmo (MED Medical Officer) Date Prepared: 16/07/2021 10:30 Signature: __OM__
---	---

The information enclosed (and any accompanying documents) is confidential and may be legally privileged. If the reader of this document is not the intended recipient you are notified that any use, disclosure, copying or distribution of the information is prohibited. If you receive this document/information in error please contact the issuing facility immediately.

Date printed: 16/7/2021 14:55

Patient Medication Summary
Pharmacy Department, My Hospital
For Patient EXAMPLE

Date: 16/07/2021

MRN: 1234567

PRODUCT DESCRIPTION	BREAKFAST	LUNCH	TEA	BEDTIME	COMMENTS
CYCLOSPORIN (NEORAL) 100MG CAPSULES	1		1		Unchanged. To prevent graft or organ rejection.
MYCOPHENALATE (CELLCEPT) 250MG CAPSULES	3		3		Unchanged. To prevent graft or organ rejection. Do not crush tablets. Can be taken with or without food.
GLICLAZIDE MR (APO-GLICLAZIDE MR) 30MG TABLETS	1		1		Unchanged. To control blood sugar. Take with or immediately after food.
WARFARIN (COUMADIN) 1MG, 2MG, 5MG TABLETS			See 'comments'		Dose changed. Current dose at 16/10/2013 is 5.5 mg in the evening. Dose to be adjusted by GP on 21/10/2013.
PARACETAMOL (PANADOL OSTEO) 665MG TABLETS	2	2	2		Unchanged. To reduce pain.
SIMVASTATIN (LIPEX) 80MG TABLETS				1	Unchanged. To lower cholesterol
The following medicine was CEASED. Do not take this medicine without further notice: AMLODOPINE (NORVASC) 5mg					CEASED as causing dizziness. GP to review need at next visit.
No known allergies or adverse drug reactions.					

Please bring this chart with you to each visit to your Doctor, Pharmacist, Dentist or other Health Care Professional

Prepared by: Sue Smith